

1 **Association of glomerular hyperfiltration with serum chemokine levels and**
2 **metabolic features in prepubertal children with overweight/obesity**

3 María L. Muzzio a,b, María L. Kabakian c, Yanina Morosán-Allo d, Silvia Ferrari e, Poupak Fallahi
4 e, Jorgelina Fernández c, María P. Santucci c, Cristina Andrés-Lacueva f,g, Alessandro Antonelli e,
5 Gabriela Brenta d, Tomás Meroño a,b,f,*

6 *a Depto. de Bioquímica Clínica, Facultad de Farmacia y Bioquímica, Universidad de Buenos Aires,*
7 *Buenos Aires, Argentina*

8 *b Laboratorio Central, Complejo Médico Churruca-Visca, Buenos Aires, Argentina*

9 *c Servicio de Diabetes y Nutrición Infanto-Juvenil, Complejo Médico Churruca-Visca, Buenos Aires,*
10 *Argentina*

11 *d Servicio de Endocrinología y Metabolismo, Unidad Asistencial “Dr. César Milstein”, Buenos*
12 *Aires, Argentina*

13 *e Department of Clinical and Experimental Medicine, University of Pisa, Pisa, Italy*

14 *f Biomarkers and Nutrimetabolomics Laboratory, Department of Nutrition, Food Sciences and*
15 *Gastronomy, Food Technology Reference Net (XaRTA), Nutrition and Food Safety Research Institute*
16 *(INSA), Faculty of Pharmacy and Food Sciences, University of Barcelona, 08028 Barcelona, Spain*

17 *g CIBER de Fragilidad y Envejecimiento Saludable (CIBERfes), Instituto de Salud Carlos III, 08028*
18 *Barcelona, Spain*

19 **KEYWORDS**

20 *Obesity;*

21 *Children;*

22 *Chemokines;*

23 *Hyperfiltration;*

24 *Renal disease*

25 * Corresponding author. Biomarkers and Nutrimetabolomics Laboratory, Department of Nutrition,
26 Food Sciences and Gastronomy, Faculty of Pharmacy and Food Sciences, University of Barcelona,
27 Av. de Joan XXIII, 27-31, 08028 Barcelona, Spain. E-mail address: tomasmero@yahoo.com.ar (T.
28 Meroño).

29 **ABSTRACT**

30 *Background and aims:* Glomerular hyperfiltration (GH) is proposed as one of the earliest events in
31 obesity (OB)-associated renal disease. Children with GH and type-1 diabetes showed increased
32 chemokine levels. Chemokine associations with glomerular filtration rate (GFR) and metabolic
33 features in prepubertal children with overweight (OW)/OB are unknown. *Methods and results:* Cross-
34 sectional study. 75 prepubertal children (aged: 9.0 ± 1.7 years) with OW/OB were studied. Clinical
35 and metabolic characteristics (including non-esterified fatty acids, NEFA) and GFR (combined
36 Zappitelli equation) were assessed. GH was defined as $GFR > 135 \text{ ml/min.1.73 m}^2$. Serum levels of
37 regulated on activation, normal T cell expressed and secreted (RANTES)/CCL5, interleukin-8 (IL-
38 8)/CXCL8 and monokine-induced by interferon- γ (MIG)/CXCL9 were measured by ELISA. Age-
39 and sex-adjusted correlations and differences were tested. 48% of the cohort was female and 13%
40 were OW, 54% OB and 33% severe OB. Prepubertal children with GH showed lower z-BMI (-12%),
41 NEFA (-26%) and uric acid (-22%) than those without GH (all $p < 0.05$). Similarly to high sensitivity
42 C-reactive protein (hsCRP), there were no differences in serum chemokines between children with
43 GH or not (all $p > 0.05$). Adjusted correlations were significant for RANTES and z-BMI ($r = 0.26$; p
44 < 0.05) and for MIG with z-BMI ($r = -0.26$; $p < 0.05$) and with NEFA ($r = 0.27$; $p < 0.05$). *Conclusion:*
45 GH was not associated with higher chemokine levels in prepubertal children with OW/OB. Decreased
46 rather than elevated GFR values were correlated with obesity and worse metabolic profiles.
47 Chemokines levels in children with severe OB suggest a regulation of the immune response. Follow-
48 up studies are needed to address the clinical implications of these findings.

49 **Introduction**

50 Childhood obesity (OB) is a growing health problem worldwide [1]. Children with overweight (OW)
51 and/or OB are at increased risk for developing several comorbidities during adult age such as type 2
52 diabetes and renal and cardiovascular diseases, among others [2,3]. In the development of obesity-
53 associated renal disease an early stage of elevated glomerular filtration rate (GFR) or glomerular
54 hyperfiltration (GH) is proposed [4]. Whether elevated GFR is a relevant clinical sign indicating early
55 impairment in renal function in children with OB is a matter of debate [5e9]. Some reports described

56 increased GFR in children with OB in comparison with normal-weight children; however, others
57 showed decreased GFR levels [5e9]. In part, controversy arises from the difficulties associated with
58 the definition of GH. Some authors used an estimated GFR calculated by the modified Schwartz
59 equation, which uses only serum creatinine concentration, while others used equations with cystatin
60 C concentration alone or with a combination of both creatinine and cystatin C levels [5,7,8]. Also,
61 different authors employed different cut off points of GFR, although a value $>135 \text{ ml/min.1.73 m}^2$ is
62 frequently used in the literature [5,10]. In adults with OB and obesity related glomerulopathy (ORG),
63 GH is commonly associated with metabolic alterations and increased inflammatory markers [4].
64 Whether such associations are also found in prepubertal children with OB have not been fully
65 described. Lee et al. [11] showed increased triglycerides and markers of insulin resistance in
66 adolescents with GH. In addition, Di Bonito et al. [12] showed that OB children (6e16 years) with an
67 altered GFR (defined as $>80\text{th}$ percentile or $<20\text{th}$ percentile) were at increased risk of presenting
68 elevated blood pressure (BP), impaired fasting glucose (IFG) and elevated white blood cell counts.
69 Non-esterified fatty acids (NEFA) are at the hallmark of metabolic abnormalities and endothelial
70 dysfunction in adults with OB and insulin resistance [13]. Impaired lipolysis and elevated NEFA in
71 adults with OB were linked to ectopic fat deposition in several organs, including the kidneys [14].
72 “Fatty” kidney could be one possible explanation to the impairment of renal function in OB [14].
73 Indeed, a recent study showed that single nucleotide polymorphisms (SNPs) predisposing to ectopic
74 fat accumulation were associated with lower GFR values in OB children with non-alcoholic fatty
75 liver disease [15]. NEFA levels and its association with GFR and inflammatory markers in
76 prepubertal children has been scarcely studied. A link between inflammatory markers and GH was
77 suggested by a positive correlation between GFR and myeloperoxidase activity in a Portuguese cohort
78 of prepubertal children with OW/OB [16]. However, most of the studies regarding GH and
79 chemokines were carried out in children and adolescents with type 1 diabetes. In particular, children
80 with type 1 diabetes and GH showed elevated urinary excretion of monocyte chemoattractant protein
81 (MCP)-1, regulated on activation, normal T cell expressed and secreted (RANTES) and eotaxin in
82 comparison with those without GH [17]. T lymphocyte chemokines, like RANTES/CCL5 and
83 interleukin-8 (IL-8)/ CXCL8, are secreted by adipose tissue and are proposed as important mediators

84 in the cross-talk between OB and cardiovascular disease [18e20]. Another chemokine relevant to
85 vascular health is the monokine induced by interferon- γ (MIG)/CXCL9 which is postulated as an
86 important mediator in hypertension and cardiovascular disease [21]. In children with OB, levels of
87 RANTES, IL-8 and MIG decreased with body weight reductions or BP changes during intervention
88 studies [22,23]. Then, these chemokines may be involved in early changes of the GFR in prepubertal
89 children with OB. The aim of the present study was to characterize the correlations of serum
90 chemokines with GFR and metabolic alterations in a cohort of prepubertal children with OW/ OB. A
91 secondary analysis was performed to explore differences between children with or without severe
92 OB.

93 **Methods**

94 **Sample size, study design and population**

95 Sample size was calculated in the base of a previous study on correlations between chemokines and
96 GFR in children and adolescents [17]. As it was done in pediatric patients with type 1 diabetes, a-
97 priori we multiplied the result by 1.5 for final sample size estimation. The number of patients to be
98 studied with an effect size of 0.381, 80% power and two-tailed significance level of 0.05, multiplied
99 by 1.5 was 73. The studied cohort was part from an ongoing prospective study which consecutively
100 recruited every child between 5 and 17 years with OW or OB who consulted to the Pediatric Diabetes
101 and Nutrition Service of the Complejo Médico Churruca Visca, Buenos Aires, Argentina, during
102 April 2017 to October 2018. Out of the 183 patients recruited, we filtered the prepubertal ones
103 (n=109). From these, a simple random sample of 80 patients was used for the analysis. At the time of
104 measuring high sensitivity C reactive protein (hsCRP), 5 patients had to be excluded in the basis of a
105 value >10 mg/l. The final number of prepubertal children included in this cross-sectional study was
106 75. All patients, including the group of normal weight children, and their parents or responsible adult
107 signed the assent and consent forms, respectively, before participation in the study. The protocol was
108 approved by the Ethics Committee of the Complejo Médico Churruca-Visca. The general exclusion
109 criteria for the study were diabetes, hypothyroidism or treatment with levothyroxine, concomitant

110 autoimmune or liver pathologies, an urinary albumin-creatinine ratio >30 mg/g, clinical signs of an
111 infectious disease or a hsCRP >10 mg/l.

112 **Clinical and laboratory measurements**

113 All the patients underwent a clinical interview in the presence of their parents or responsible adult in
114 which height and weight were determined without shoes and wearing light clothes. Body mass index
115 (BMI) was calculated and BMI classification was performed according to the z-score from the World
116 Health Organization (WHO) [24]. Severe OB was defined by a z-BMI > 3.0. BP was measured
117 twice after a 5 min rest in the sitting position and the average of both measurements was recorded.
118 The American Academy of Pediatrics (AAP) 2017 guidelines were used to calculate BP z-scores and
119 to define elevated BP [25]. Waist circumference (WC) was measured at the mid-point between the
120 iliac crest and the 10th rib and elevated WC classified as >90th percentile of the Bogalusa Heart
121 Study [26]. After a 12 h overnight fast, blood samples were drawn and general lab measurements
122 were carried out within a 24 h period. Serum samples were frozen at -70°C for the hsCRP and
123 chemokines measurements. General laboratory tests were carried out in a Unicel DxC 800 (Beckman
124 Coulter, USA) autoanalyzer by standardized methods. Cystatin C concentration was measured with
125 an immunoturbidimetric assay (Diasyme, USA) and NEFA by a colorimetric assay (Randox, UK)
126 according to manufacturer indications. HsCRP was measured by a nephelometric assay (Image,
127 Beckman Coulter, USA). GFR was estimated by the combined creatinine and cystatin C Zappitelli
128 formula and GH was defined in presence of a GFR >135 ml/min.1.73 m² [27]. In order to verify the
129 GH cut-off suggested by bibliography [28,29], 26 samples of prepubertal normal weight children
130 were tested for creatinine and cystatin C and only one patient showed an GFR outside the defined
131 reference range (90-135 ml/min.1.73 m²). These 26 children (44% female, aged: 9-12 years) were
132 attended in the same Hospital service as the patients and, besides being normal-weight [z-BMI:
133 median (Q1-Q3), 0.07 (-0.67-0.62)] met the same exclusion criteria as the patients with OW/OB.
134 The range of GFR from these children was between 98 and 136 ml/min.1.73 m², with a median at
135 117 ml/min.1.73 m² and Q1-Q3 111-126 ml/min.1.73 m². Serum levels of RANTES/CCL5,
136 MIG/CXCL9 and IL-8/CXCL8 were measured by ELISA (R&D systems, USA). Cardiometabolic

137 risk factors were categorized according to the definition of metabolic syndrome of Cook et al. [30]
138 modified by the Sociedad Argentina de Pediatría. The following cut-offs were used: abdominal OB,
139 WC > 90th percentile; elevated BP, systolic or diastolic BP > 90th percentile; impaired fasting
140 glucose (IFG), glucose > 100 mg/dl; hypertriglyceridemia (HTG), TG > 110 mg/dl and low HDL-C,
141 HDL-C < 40 mg/dl.

142 **Statistical analyses**

143 Variables were tested for normality by the Shapiro-Wilks test. Normally distributed variables are
144 shown as mean \pm standard deviation (sd) and skewed distributed variables as median (Q1eQ3).
145 According to data distribution, Pearson or Spearman correlation tests were used. The different groups
146 were compared by Student T test or Mann-Whitney U test for normally or skewed distributed
147 variables, respectively. Following log-transformation of skewed variables partial correlation test or
148 ANCOVA was used to adjust by age and sex. Residuals of multivariate tests were checked for
149 normality to evaluate the models. Categorical variables were compared with the Fisher Exact test.
150 SPSS 25.0 (IBM, USA) was used for statistical analyses. Tests were considered statistically
151 significant with a significance value (p) < 0.05.

152 **Results**

153 **Population characteristics and evaluation of renal function**

154 The studied population comprised 75 prepubertal children (48% female) of which 13% showed OW,
155 54% OB and 33% severe OB. The overall prevalence of metabolic syndrome components were:
156 abdominal OB: 73%, low HDL-C: 57%, HTG: 39%, elevated BP: 23%, and IFG: 9%. Metabolic
157 syndrome was diagnosed in 32% of the children and GH was found in 16%. Clinical and laboratory
158 characteristics according to the presence of GH are shown in Table 1. Prepubertal children with GH
159 showed lower z-BMI, NEFA and uric acid than those with normal GFR. GFR correlated with age (r
160 Z -0.32; p Z 0.005), NEFA (r Z -0.30; p Z 0.011) and uric acid levels (r Z -0.29; p Z 0.011). These
161 correlation were independent of sex and age and sex, accordingly. Besides its inverse correlation with
162 GFR, uric acid levels correlated with the number of cardiometabolic risk factors (r Z 0.37; p < 0.001),

LDL-C ($r = 0.30$; $p = 0.010$), TG/ HDL-C ($r = 0.28$; $p = 0.015$), NEFA ($r = 0.39$; $p < 0.001$) and hsCRP ($r = 0.29$; $p = 0.014$). These correlations, except that with TG/HDL-C, remained statistically significant when adjusting by age and sex.

Serum chemokine levels and its correlation with renal function and metabolic characteristics

Similarly to the inflammatory marker hsCRP (Table 1), there were no differences in serum chemokines between children with GH or not [RANTES: 27.2 (22.4e29.0) vs. 26.9 (23.6e30.0) ng/ml; MIG: 420 (329e523) vs. 454 (276e686) pg/ml; and IL-8: 551 (311e782) vs. 630 (406e922) pg/ml, respectively, all $p > 0.05$]. MIG and IL-8 were both inversely correlated with GFR ($r = -0.27$, $p = 0.035$; and $r = -0.25$, $p = 0.022$, respectively). These correlations were no longer significant when tests were adjusted by age and sex ($p > 0.05$). RANTES levels were not correlated with GFR ($p > 0.05$). Correlations of serum chemokine levels with z-BMI and NEFA are shown in Fig. 1. Strikingly, while RANTES was directly correlated with z-BMI, MIG was inversely correlated with z-BMI and positively with NEFA (Fig. 1). IL-8 levels were closely correlated with MIG levels ($r = 0.78$; $p < 0.001$), and IL-8 correlations with z-BMI and NEFA mirrored those of MIG (Fig. 1). In analyses adjusted by age and sex, only RANTES correlation with z-BMI and both MIG correlations remained statistically significant (all $p < 0.05$). None of the other metabolic characteristics were correlated with the serum chemokine levels. HsCRP levels were significantly correlated with z-BMI ($r = 0.45$; $p < 0.001$) and this correlation remained significant even when adjusting by age and sex ($p < 0.05$).

Subgroup analysis according to severe obesity

Prepubertal children with severe OB were younger and showed a worse cardiometabolic profile (Table 2). Strikingly, children with severe OB did not present GH which was exclusively seen in children without severe OB (Table 1). Children with severe OB presented higher levels of hsCRP but lower levels of the chemokines IL-8 and MIG. Differences in hsCRP and MIG remained statistically significant in models adjusted by age and sex (Table 2). Adjusted differences were also significant for HOMA-IR and TG.

Discussion

189 The present study shows that serum chemokines concentrations were not different between
190 prepubertal children with or without GH. Moreover, children with GH were less obese and did not
191 showed higher levels of cardiometabolic risk factors. In fact, children with severe OB did not show
192 a higher GFR, in spite of being younger. Altogether, these results suggest the existence of a pool of
193 OB children with an early decline of GFR in which GH could have happened at even younger ages,
194 as declining GFR values in childhood OB were also noticed by others [9,15]. As an alternative, the
195 present definition of GH ($GFR > 135 \text{ ml/min.1.73 m}^2$) might be limited to detect early renal
196 impairment. Whatever the case, this insight is clinically relevant as prepubertal children with OW/OB
197 developing early alterations in renal function would not be identified by current biochemical markers.
198 In this cohort of prepubertal children there were no correlations between GFR and z-BMI. Moreover,
199 instead of positive, negative correlations of GFR with NEFA and uric acid were seen. Both elevated
200 NEFA and uric acid are not only markers but also proposed mediators of cardiometabolic
201 derangements in OB. Uric acid was positively correlated with cardiometabolic risk factors, the
202 TG/HDL-C ratio, LDL-C, NEFA and hsCRP, similarly to what it was reported by other studies in
203 prepubertal children with OB [31]. Thus, our data supports the notion that OB and its metabolic
204 alterations could lead to a reduction of GFR rather than to an increase in prepubertal children. This
205 result contrasts with the studies that used different GFR equations, like Bouvet or modified Schwartz
206 [5,9,27]. In particular, Bouvet equation includes weight as a variable and then higher but not lower
207 GFR values in OB children could be expected [5]. On the other hand, the use of Schwartz equation
208 could lead to bias related to body distribution of fat and lean mass and its influence on creatinine
209 levels [9]. Although other sources of variation in serum cystatin C levels, like increased secretion by
210 adipose tissue cannot be dismissed [32], our results support those of a Portuguese cohort of 313
211 prepubertal children (48% with OW/OB) [7]. In the latter, not only GFR using Zappitelli equation
212 was negatively correlated with z-BMI, but also 24h creatinine clearance [7]. Nonetheless, the impact
213 of OB over GFR in childhood remains to be determined in longitudinal studies. Prepubertal children
214 with severe OB showed a worse cardiometabolic profile and, paradoxically, elevated hsCRP and
215 decreased MIG levels. In this point, it should be noticed that while hsCRP were positively correlated
216 with z-BMI, MIG did in an opposite way. Then, the IL-1b d IL-6 d CRP axis could reflect the

217 developing inflammatory state set up by the onset and progression of fat accumulation, as it was also
218 noticed in cross-sectional and intervention studies involving prepubertal children [33e36]. On the
219 other hand, the inverse correlation between z-BMI and MIG could be related to an adaptative
220 counterregulation involving the adipose tissue regulatory T cells (Treg) [37]. This adaptative Treg
221 response was increased in OB vs. lean humans and was more relevant in subcutaneous than visceral
222 adipose tissue [38,39]. Such specific responses in fat depots is relevant to the interpretation of data
223 on prepubertal children with OB. Before puberty, gain of subcutaneous fat mass is more relevant,
224 while from puberty onset and onward visceral fat accumulation is more significant [40,41]. Indeed,
225 when comparing prepubertal children with or without OB differences in subcutaneous adipose tissue
226 were quantitatively more significant than those in visceral adipose tissue [36,42]. Also, subcutaneous
227 rather than visceral adipose tissue was correlated with reduced insulin sensitivity and ectopic fat
228 accumulation before puberty [36,42,43]. On the other hand, MIG negative correlation with NEFA
229 could be at some point attributed to the known effects of IFN-g on lipolysis [44]. In regards to IL-8,
230 its correlations with metabolic characteristics resembled those of MIG and an inverse correlation
231 between IL-8 and z-BMI was also reported in a cohort of prepubertal children [42]. Although IL-8
232 expression was demonstrated in adipose tissue, the correlation between MIG and IL-8 was
233 independent of z-BMI in our study (data not shown). Thus, another source of MIG and IL-8 (like
234 monocytes) could be suspected in prepubertal children with OB. Indeed, IFN-g promoted the
235 secretion of IL-8 and MIG by monocytes in synergy with tumor necrosis factor (TNF)-a [45]. The
236 positive correlation between RANTES and z-BMI is in line with T cell infiltration of adipose tissue
237 as a consequence of fat mass expansion [38]. T cell infiltration is proposed as one of the earliest
238 events in OB [46]. Thus, elevation of RANTES could be related to an adaptative response to OB in
239 children, as RANTES exhibited antiapoptotic activities in adipose tissue macrophages [47].
240 Nonetheless, elevated RANTES levels were correlated with arterial stiffness, an early marker of
241 endothelial dysfunction, in healthy prepubertal children [48]. Then, RANTES levels could mirror
242 maladaptative responses in OB and it might be related with its cardiometabolic consequences. The
243 cross-sectional design of our study did not allow us to draw conclusions on mechanisms and our
244 hypothesis to explain the correlations observed are speculative. Nonetheless, few studies focused in

the relationship between chemokines, NEFA and GFR in prepubertal children with OB. Moreover, children with severe OB represent a group of understudied individuals, especially before pubertal development. In the present cohort, severe OB was present in 33% of the children with z-BMI values ranging: 3.03e6.72. The evaluation of these patients by separate gave further support to the notion of declining GFR (and increasing uric acid levels) in children with OB as possible early events in the development of renal disease. Future studies will address if control of uric acid levels may help preserve renal function in childhood OB. On the other hand, the relatively low number of patients included, could be considered a shortcoming and raises the possibility to have missed relevant results. Among the strengths of the study there are two to be highlighted: a) the exclusion of patients with clinical or biochemical signs (hsCRP > 10 mg/l) of infectious or inflammatory conditions that minimized the possibility of bias in chemokine levels due to concomitant pathological processes and b) the use of sex and age-adjusted analysis to draw the main conclusions of the study. In conclusion, in prepubertal children with OW/OB serum levels of RANTES, IL-8 and MIG were not correlated with the GFR. Increasing z-BMI and worse metabolic profiles were associated with decreased rather than elevated GFR values. While RANTES was positively correlated with z-BMI, MIG showed an inverse correlation. Follow-up studies are needed to address the clinical implications of these findings.

Acknowledgments

This study could not have been done without the collaboration of the technicians of the Laboratorio Central of the Complejo Médico Churruca-Visca, Buenos Aires, Argentina. The present work was partially funded by a grant from Fundación Alberto J. Roemmers. TM would like to thank the “Juan de la Cierva” program from MINECO (IJCI-2017- 32534). The authors have no conflict of interest to declare.

References

[1] World Health Organization. WHO | Facts and figures on childhood obesity. WHO; 2019.

270 [2] Vivante A, Golan E, Tzur D, Leiba A, Tirosh A, Skorecki K, et al. Body mass index in 1.2 million
 271 adolescents and risk for end-stage renal disease. *Arch Intern Med* 2012;172:1644e50. <https://doi.org/10.1001/2013.jamainternmed.85>.
 272

273 [3] Buscot M-J, Thomson RJ, Juonala M, Sabin MA, Burgner DP, Lehtimäki T, et al. Distinct child-
 274 to-adult body mass index trajectories are associated with different levels of adult cardiometabolic
 275 risk. *Eur Heart J* 2018;39:2263e70. <https://doi.org/10.1093/eurheartj/ehy161>.
 276

276 [4] D'Agati VD, Chagnac A, de Vries APJ, Levi M, Porrini E, Herman- Edelstein M, et al. Obesity-
 277 related glomerulopathy: clinical and pathologic characteristics and pathogenesis. *Nat Rev Nephrol*
 278 2016; 12:453e71. <https://doi.org/10.1038/nrneph.2016.75>.
 279

279 [5] Franchini S, Savino A, Marcovecchio ML, Tumini S, Chiarelli F, Mohn A. The effect of obesity
 280 and type 1 diabetes on renal function in children and adolescents. *Pediatr Diabetes* 2015;16:427e33.
 281 <https://doi.org/10.1111/pedi.12196>.
 282

282 [6] van Dam M, Rijks J, Dorenbos E, Horuz F, van Dael K, Vreugdenhil A. The effect of one year
 283 lifestyle intervention on eGFR in children and adolescents with overweight, obesity and morbid
 284 obesity. *Sci Rep* 2019;9:4504. <https://doi.org/10.1038/s41598-019-40767-4>.
 285

285 [7] Correia-Costa L, Afonso AC, Schaefer F, Guimarães JT, Bustorff M, Guerra A, et al. Decreased
 286 renal function in overweight and obese prepubertal children. *Pediatr Res* 2015;78:436e44. <https://doi.org/10.1038/pr.2015.130>.
 287

288 [8] Önerli Salman D, S, ıklar Z, Çullas İlarslan EN, Özçakar ZB, Kocaay P, Berberoğlu M.
 289 Evaluation of renal function in obese children and adolescents using serum cystatin C levels,
 290 estimated glomerular filtration rate formulae and proteinuria: which is most useful? *J Clin Res Pediatr*
 291 *Endocrinol* 2019;11:46e54. <https://doi.org/10.4274/jcrpe.galenos.2018.2018.0046>.
 292

292 [9] Marzuillo P, Grandone A, Di Sessa A, Guarino S, Diplomatico M, Umano GR, et al.
 293 Anthropometric and biochemical determinants of estimated glomerular filtration rate in a large cohort
 294 of obese children. *J Ren Nutr* 2018;28:359e62. <https://doi.org/10.1053/j.jrn.2018.01.001>.

295 [10] Karnebeek K, Thapar S, Willeboordse M, van Schayck OCP, Vreugdenhil ACE. Comorbidities
 296 in primary vs secondary school children with obesity and responsiveness to lifestyle intervention. *J*
 297 *Clin Endocrinol Metab* 2019;104:3803e11. <https://doi.org/10.1210/jc.2018-02318>.

298 [11] Lee AM, Charlton JR, Carmody JB, Gurka MJ, DeBoer MD. Metabolic risk factors in
 299 nondiabetic adolescents with glomerular hyperfiltration. *Nephrol Dial Transplant* 2016;32:gfw231.
 300 <https://doi.org/10.1093/ndt/gfw231>.

301 [12] Di Bonito P, Sanguigno E, Forziato C, Di Fraia T, Moio N, Cavuto L, et al. Glomerular filtration
 302 rate and cardiometabolic risk in an outpatient pediatric population with high prevalence of obesity.
 303 *Obesity* 2014;22:585e9. <https://doi.org/10.1002/oby.20497>.

304 [13] Imrie H, Abbas A, Kearney M. Insulin resistance, lipotoxicity and endothelial dysfunction.
 305 *Biochim Biophys Acta Mol Cell Biol Lipids* 2010;1801:320e6.
 306 <https://doi.org/10.1016/j.bbalip.2009.09.025>.

307 [14] Gai Z, Wang T, Visentin M, Kullak-Ublick G, Fu X, Wang Z. Lipid accumulation and chronic
 308 kidney disease. *Nutrients* 2019;11:722. <https://doi.org/10.3390/nu11040722>.

309 [15] Marzuillo P, Di Sessa A, Guarino S, Capalbo D, Umamo GR, Pedullà M, et al. Nonalcoholic
 310 fatty liver disease and eGFR levels could be linked by the PNPLA3 I148M polymorphism in children
 311 with obesity. *Pediatr Obes* 2019;14:e12539. <https://doi.org/10.1111/ijpo.12539>.

312 [16] Correia-Costa L, Sousa T, Morato M, Cosme D, Afonso J, Moura C, et al. Association of
 313 myeloperoxidase levels with cardiometabolic factors and renal function in prepubertal children. *Eur*
 314 *J Clin Invest* 2016;46:50e9. <https://doi.org/10.1111/eci.12564>.

315 [17] Har R, Scholey JW, Daneman D, Mahmud FH, Dekker R, Lai V, et al. The effect of renal
 316 hyperfiltration on urinary inflammatory cytokines/ chemokines in patients with uncomplicated type
 317 1 diabetes mellitus. *Diabetologia* 2013;56:1166e73. <https://doi.org/10.1007/s00125-013-2857-5>.

318 [18] Wu H, Ghosh S, Perrard XD, Feng L, Garcia GE, Perrard JL, et al. Tcell accumulation and
319 regulated on activation, normal T cell expressed and secreted upregulation in adipose tissue in
320 obesity. *Circulation* 2007;115:1029e38. <https://doi.org/10.1161/CIRCULATIONAHA.106.638379>.

321 [19] Kim C-S, Park H-S, Kawada T, Kim J-H, Lim D, Hubbard NE, et al. Circulating levels of MCP-
322 1 and IL-8 are elevated in human obese subjects and associated with obesity-related parameters. *Int*
323 *J Obes* 2006;30:1347e55. <https://doi.org/10.1038/sj.ijo.0803259>.

324 [20] Hoch M, Eberle AN, Peterli R, Peters T, Seboek D, Keller U, et al. LPS induces interleukin-6
325 and interleukin-8 but not tumor necrosis factor-a in human adipocytes. *Cytokine* 2008;41:29e37.
326 <https://doi.org/10.1016/j.cyto.2007.10.008>.

327 [21] Youn J-C, Yu HT, Lim BJ, Koh MJ, Lee J, Chang D-Y, et al. Immunosenescent CD8 β T cells
328 and C-X-C chemokine receptor type 3 chemokines are increased in human hypertension.
329 *Hypertension* 2013;62:126e33. <https://doi.org/10.1161/HYPERTENSIONAHA.113.00689>.

330 [22] Rubin DA, Duran AT, Haqq AM, Gertz ER, Dumont-Driscoll M. Changes in cardiometabolic
331 markers in children with Prader-Willi syndrome and nonsyndromic obesity following participation
332 in a home-based physical activity intervention. *Pediatr Obes* 2018;13: 734e43.
333 <https://doi.org/10.1111/ijpo.12462>.

334 [23] Övünç Hacıhamdioğlu D, Zeybek C, Gök F, Pekel A, Musabak U. Elevated urinary T helper
335 1 chemokine levels in newly diagnosed hypertensive obese children. *J Clin Res Pediatr Endocrinol*
336 2015;7: 175e82. <https://doi.org/10.4274/jcrpe.1917>.

337 [24] de Onis M. Development of a WHO growth reference for schoolaged children and adolescents.
338 *Bull World Health Organ* 2007; 85:660e7. <https://doi.org/10.2471/BLT.07.043497>.

339 [25] Flynn JT, Kaelber DC, Baker-Smith CM, Blowey D, Carroll AE, Daniels SR, et al. Clinical
340 practice guideline for screening and management of high blood pressure in children and adolescents.
341 *Pediatrics* 2017;140:e20171904. <https://doi.org/10.1542/peds.2017-1904>.

342 [26] Freedman DS, Serdula MK, Srinivasan SR, Berenson GS. Relation of circumferences and
343 skinfold thicknesses to lipid and insulin concentrations in children and adolescents: the Bogalusa
344 Heart Study. *Am J Clin Nutr* 1999;69:308e17. <https://doi.org/10.1093/ajcn/69.2.308>.

345 [27] Bacchetta J, Cochat P, Rognant N, Ranchin B, Hadj-Aissa A, Dubourg L. Which creatinine and
346 cystatin C equations can be reliably used in children? *Clin J Am Soc Nephrol* 2011;6:552e60.
347 <https://doi.org/10.2215/CJN.04180510>.

348 [28] Fadrowski JJ, Neu AM, Schwartz GJ, Furth SL. Pediatric GFR estimating equations applied to
349 adolescents in the general population. *Clin J Am Soc Nephrol* 2011;6:1427e35. <https://doi.org/10.2215/CJN.06460710>.

351 [29] Cachat F, Combescure C, Cauderay M, Girardin E, Chehade H. A systematic review of
352 glomerular hyperfiltration assessment and definition in the medical literature. *Clin J Am Soc Nephrol*
353 2015; 10:382e9. <https://doi.org/10.2215/CJN.03080314>.

354 [30] Cook S, Weitzman M, Auinger P, Nguyen M, Dietz WH. Prevalence of a metabolic syndrome
355 phenotype in adolescents: findings from the Third National Health and Nutrition Examination
356 Survey, 1988-1994. *Arch Pediatr Adolesc Med* 2003;157:821e7. <https://doi.org/10.1001/archpedi.157.8.821>.

358 [31] Valle M, Martos R, Cañete MD, Valle R, van Donkelaar EL, Bermudo F, et al. Association of
359 serum uric acid levels to inflammation biomarkers and endothelial dysfunction in obese prepubertal
360 children. *Pediatr Diabetes* 2015;16:441e7. <https://doi.org/10.1111/pedi.12199>.

361 [32] Naour N, Fellahi S, Renucci J-F, Poitou C, Rouault C, Basdevant A, et al. Potential contribution
362 of adipose tissue to elevated serum cystatin C in human obesity. *Obesity (Silver Spring)* 2009;17:
363 2121e6. <https://doi.org/10.1038/oby.2009.96>.

364 [33] Galcheva SV, Iotova VM, Yotov YT, Bernasconi S, Street ME. Circulating proinflammatory
365 peptides related to abdominal adiposity and cardiometabolic risk factors in healthy prepubertal
366 children. *Eur J Endocrinol* 2011;164:553e8. <https://doi.org/10.1530/EJE-10-1124>.

367 [34] Stoppa-Vaucher S, Dirlewanger MA, Meier CA, de Moerloose P, Reber G, Roux-Lombard P, et
368 al. Inflammatory and prothrombotic states in obese children of European descent. *Obesity* 2012;20:
369 1662e8. <https://doi.org/10.1038/oby.2012.85>.

370 [35] Martos R, Valle M, Morales RM, Cañete R, Gascón F, Urbano MM. Changes in body mass
371 index are associated with changes in inflammatory and endothelial dysfunction biomarkers in obese
372 prepubertal children after 9 months of body mass index SD score loss. *Metabolism* 2009;58:1153e60.
373 <https://doi.org/10.1016/j.metabol.2009.03.017>.

374 [36] Maffei C, Manfredi R, Trombetta M, Sordelli S, Storti M, Benuzzi T, et al. Insulin sensitivity
375 is correlated with subcutaneous but not visceral body fat in overweight and obese prepubertal
376 children. *J Clin Endocrinol Metab* 2008;93:2122e8. <https://doi.org/10.1210/jc.2007-2089>.

377 [37] Zeng Q, Sun X, Xiao L, Xie Z, Bettini M, Deng T. A unique population: adipose-resident
378 regulatory T cells. *Front Immunol* 2018;9: 2075. <https://doi.org/10.3389/fimmu.2018.02075>.

379 [38] Zeyda M, Huber J, Prager G, Stulnig TM. Inflammation correlates with markers of T-cell subsets
380 including regulatory T cells in adipose tissue from obese patients. *Obesity* 2011;19:743e8. <https://doi.org/10.1038/oby.2010.123>.

381

382 [39] Pereira S, Teixeira L, Aguilar E, Oliveira M, Savassi-Rocha A, Pelaez JN, et al. Modulation of
383 adipose tissue inflammation by FOXP3⁺ Treg cells, IL-10, and TGF- β in metabolically healthy class
384 III obese individuals. *Nutrition* 2014;30:784e90. <https://doi.org/10.1016/j.nut.2013.11.023>.

385 [40] Huang TT-K, Johnson MS, Figueroa-Colon R, Dwyer JH, Goran MI. Growth of visceral fat,
386 subcutaneous abdominal fat, and total body fat in children. *Obesity* 2001;9:283e9. <https://doi.org/10.1038/oby.2001.35>.

387

388 [41] Gyllenhammer LE, Alderete TL, Toledo-Corral CM, Weigensberg M, Goran MI. Saturation of
389 subcutaneous adipose tissue expansion and accumulation of ectopic fat associated with metabolic
390 dysfunction during late and post-pubertal growth. *Int J Obes (Lond)* 2016;40:601e6.
391 <https://doi.org/10.1038/ijo.2015.207>.

392 [42] Zabaleta J, Velasco-Gonzalez C, Estrada J, Ravussin E, Pelligrino N, Mohler MC, et al. Inverse
393 correlation of serum inflammatory markers with metabolic parameters in healthy, Black and White
394 prepubertal youth. *Int J Obes (Lond)* 2014;38:563e8. <https://doi.org/10.1038/ijo.2013.220>.

395 [43] Larson-Meyer DE, Newcomer BR, Ravussin E, Volaufova J, Bennett B, Chalew S, et al.
396 Intrahepatic and intramyocellular lipids are determinants of insulin resistance in prepubertal children.
397 *Diabetologia* 2011;54:869e75. <https://doi.org/10.1007/s00125-010-2022-3>.

398 [44] Doerrler W, Feingold KR, Grunfeld C. Cytokines induce catabolic effects in cultured adipocytes
399 by multiple mechanisms. *Cytokine* 1994;6:478e84.

400 [45] Schrier SB, Hill AS, Plana D, Lauffenburger DA. Synergistic communication between CD4⁺ T
401 cells and monocytes impacts the cytokine environment. *Sci Rep* 2016;6:34942. <https://doi.org/10.1038/srep34942>.

403 [46] Nishimura S, Manabe I, Nagasaki M, Eto K, Yamashita H, Ohsugi M, et al. CD8⁺ effector T
404 cells contribute to macrophage recruitment and adipose tissue inflammation in obesity. *Nat Med*
405 2009;15: 914e20. <https://doi.org/10.1038/nm.1964>.

406 [47] Keophiphath M, Rouault C, Divoux A, Clément K, Lacasa D. CCL5 promotes macrophage
407 recruitment and survival in human adipose tissue. *Arterioscler Thromb Vasc Biol* 2010;30:39e45.
408 <https://doi.org/10.1161/ATVBAHA.109.197442>.

409 [48] Eikendal ALM, Evelein AMV, Uiterwaal CSPM, van der Ent CK, Visseren FLJ, Bots ML, et al.
410 Relation between circulating inflammatory chemokines and vascular characteristics in healthy, young
411 children. *J Am Heart Assoc* 2015;4. <https://doi.org/10.1161/JAHA.115.002346>.




Figure 1 Correlations between serum chemokines and z-BMI and NEFA ($n = 75$). NEFA, non-esterified fatty acids; RANTES, regulated on activation, normal T cell expressed and secreted; IL, interleukin; MIG, monokine induced by interferon γ . Coefficients and significance levels shown are those of univariate tests. *Remained significant in tests adjusted by age and sex.

TABLES

Table 1 Characteristics of prepubertal children with OW/OB according to the presence of GH ($n = 75$).

	GH ($n = 12$)	Normal GFR ($n = 63$)	<i>p</i>
Age (years)	8.2 ± 1.2	9.2 ± 1.7	0.214
Female sex ($n, \%$)	8 (66)	28 (44)	0.212
z-BMI	2.5 (2.0–2.6)	2.8 (2.4–3.7)	0.016 ^a
Severe OB ($n, \%$)	0 (0)	25 (40)	0.005
z-SBP	0.51 (−0.19 — 0.87)	0.62 (−0.19 — 1.08)	0.470
z-DBP	0.32 (−0.11 — 0.57)	0.47 (−0.31 — 0.93)	0.509
Elevated BP ($n, \%$)	2 (17)	16 (25)	0.719
WC (cm)	75 ± 7	84 ± 13	0.054
Glucose (mmol/l)	5.0 ± 0.4 [90 ± 7]	5.2 ± 0.4 [93 ± 7]	0.326
HOMA-IR	2.7 (1.4–4.7)	2.2 (1.6–3.8)	0.740
TG (mmol/l)	1.1 (0.6–1.2) [97 (57–104)]	1.2 (0.8–1.6) [103 (69–140)]	0.308
TC (mmol/l)	4.5 ± 0.6 [159 ± 22]	4.7 ± 0.8 [166 ± 29]	0.403
HDL-C (mmol/l)	1.03 (0.94–1.23) [36 (33–43)]	1.09 (0.9–1.4) [38 (33–48)]	0.115
LDL-C (mmol/l)	2.9 ± 0.5 [103 ± 17]	2.9 ± 0.7 [103 ± 26]	0.996
Non-HDL-C (mmol/l)	3.5 ± 0.7 [121 ± 23]	3.6 ± 0.8 [125 ± 28]	0.641
MetS ($n, \%$)	3 (25)	21 (33)	0.741
NEFA (mmol/l)	0.69 ± 0.27	0.87 ± 0.26	0.027 ^a
Uric acid (μmol/l)	220 ± 42 [3.7 ± 0.7]	267 ± 59 [4.5 ± 1.0]	0.013 ^a
Creatinine (μmol/l)	34 (32–38) [0.38 (0.36–0.43)]	42 (38–47) [0.47 (0.43–0.53)]	0.002 ^a
Cystatin C (mg/l)	0.65 ± 0.08	0.83 ± 0.08	<0.001 ^a
GFR (ml/min.1.73 m ²)	144 ± 10	113 ± 12	<0.001 ^a
hsCRP (mg/l)	1.5 (0.4–2.9)	2.1 (1.1–5.5)	0.165

OW, overweight; OB, obese; BMI, body mass index; SBP, systolic blood pressure; DBP, diastolic blood pressure; BP, blood pressure; WC, waist circumference; TG, triglycerides; TC, total cholesterol; HDL, high density lipoprotein; LDL, low density lipoprotein; MetS, metabolic syndrome; NEFA, non-esterified fatty acids; GFR, glomerular filtration rate (calculated with combined Zappitelli equation).

^a Differences remained statistically significant in models adjusted by age and sex. Traditional units are in brackets. Variables are expressed as mean ± standard deviation or median (Q1–Q3), according to data distribution.

Table 2 Comparison of patient's characteristics with or without severe obesity.

	Severe OB (n = 25)	OW/OB (n = 50)	p
Age (years)	8.4 ± 1.6	9.4 ± 1.6	0.014
Female sex (n,%)	8 (32)	28 (56)	0.056
z-BMI	3.9 (3.6–4.4)	2.5 (2.0–2.7)	<0.001 ^a
z-SBP	0.73 (0.15–1.39)	0.55 (-0.16 – 1.03)	0.057
z-DBP	0.44 (-0.22–1.40)	0.28 (-0.31 – 0.63)	0.027
Elevated BP (n,%)	10 (40)	8 (16)	0.042
WC (cm)	86 ± 15	81 ± 11	0.215
Glucose (mmol/l)	5.0 ± 0.4 [90 ± 8]	5.1 ± 0.4 [91 ± 7]	0.740
HOMA-IR	3.4 (1.8–6.7)	2.1 (1.4–3.7)	0.013 ^a
TG (mmol/l)	1.6 (0.9–1.7) [139 (80–154)]	1.1 (0.8–1.3) [93 (68–112)]	0.010 ^a
HDL-C (mmol/l)	1.14 (0.94–1.49) [40 (33–52)]	1.06 (0.94–1.26) [37 (33–44)]	0.356
NEFA (mmol/l)	0.78 ± 0.19	0.87 ± 0.30	0.119
MetS (n,%)	13 (52)	11 (22)	0.017
Uric acid (μmol/l)	267 ± 65 [4.5 ± 1.1]	256 ± 59 [4.3 ± 1.0]	0.416
Creatinine (μmol/l)	40 (35–42) [0.45 (0.40–0.48)]	42 (36–47) [0.47 (0.41–0.53)]	0.235
Cystatin C (mg/l)	0.80 ± 0.07	0.80 ± 0.12	0.941
GFR (ml/min.1.73 m ²)	119 ± 9	118 ± 19	0.758
hsCRP (mg/l)	4.2 (1.5–6.3)	1.5 (0.5–3.8)	0.002 ^a
RANTES (ng/ml)	28.6 (24.2–31.4)	26.3 (23.7–29.1)	0.081
IL-8 (pg/ml)	438 (352–728)	640 (488–932)	0.045
MIG (pg/ml)	294 (220–536)	526 (373–693)	0.002 ^a

OW, overweight; OB, obese; BMI, body mass index; SBP, systolic blood pressure; DBP, diastolic blood pressure; BP, blood pressure; WC, waist circumference; TG, triglycerides; HDL, high density lipoprotein; NEFA, non-esterified fatty acids; MetS, metabolic syndrome; GFR, glomerular filtration rate (calculated with combined Zappitelli equation); RANTES, regulated on activation, normal T cell expressed and secreted; IL, interleukin; MIG, monokine induced by interferon γ .

^a Differences remained statistically significant in models adjusted by age and sex. Traditional units are in brackets. Variables are expressed as mean ± standard deviation or median (Q1–Q3), according to data distribution.

